

Approach to managing Breathlessness in a cancer patient



Patient with cancer is breathless

Before planning further, determine;

1. Reversible contributors
2. mMRC Grade
3. Position on the disease trajectory
4. Prognosis (below)

Cancer-related lung
infiltration, lymphangitis
carcinomatosis, pleural
effusion, airway/SVC
obstruction, phrenic N
palsy, ascitis, severe pain.

Treatment related
lobectomy,
pneumonectomy, RT
pneumonitis, fibrosis,
blocked tracheostomy,
Pulmonary embolism


C-morbidities
Late stage COPD,
Anemia

Debility
related severe
weakness
/fatigue,
uncommon
infections

Psychological
Fear, extreme
anxiety, distress

- Correct the correctable - analgesia, tap effusions, antibiotics, broncho-dilators, inhaled / nebulised cortico-steroids, anti-depressants, anxiolytics – **activate appropriate general medicine / surgical packages**
- In selected patients - anticoagulation, blood transfusion, brachytherapy, stenting, laser, rehabilitation - **activate appropriate general medicine / surgical / oncology packages**
- In all patients – **Included in this palliative care package**
 - Non-abandonment, empathetic communications, reassurance and emotional support
 - Tripod sitting position, loosen clothes, Pursed lip breathing, use fan/ open windows, direct airflow across the face, wet the face, mindful belly breathing
 - Anxiety reduction, relaxation, distraction, visualisation, facilitate development of individualised relaxation tool-kit (accessible contents for sensory/intellectual distraction - soothing audio/ music, soft toy, cookies/ chocolates, visuals, poems / quotes, etc.)
 - Introduce the concept of Advance care (living will), maintain dialogue, clarify & document decisions, update patient record

Palliative Care Approach to managing Breathlessness in a cancer patient continued.



Medications (A) – (Prescribe medications for baseline, incident and crisis dyspnoea management)

1. Bronchodilators
 - Salbutamol 2.5-5mg.QDS via nebuliser, or 2 puffs four times per day using spacer
 - Ipratropium 250-500 micrograms up to QDS via nebuliser, or 2 puffs. via spacer device
2. Opioids¹ (American college of Chest Physicians & American Thoracic Society & NCCN – Grade 1 A recommendation)
 - Morphine 2.5mg PO 4-hourly initially, immediate release
 - Sustained release and continuous infusion are also beneficial.
 - Impact on pO₂, pCO₂, sPO₂ not clinically significant
 - Titrate to effect. 30 mg/D in opioid-naïve patients. 25% escalation of Dose for those already on Morphine for Pain Relief
 - Manage side-effects prophylactically e.g. stimulant laxative
 - Caution should be used in the elderly or in the presence of renal impairment
2. Corticosteroids – when multiple lung metastases and in lymphangitis carcinomatosa to reduce peri-tumour oedema.
 - Dexamethasone - 4-8mg OM, for a one week trial and if there is no improvement, stop.
 - Gastric mucosal protection not indicated for 5 day's steroid trial
3. Benzodiazepines - for accompanying anxiety, panic
 - Lorazepam 0.5-1mg SL. PRN 6-8hrly
 - Diazepam 2-5 mg
 - Midazolam 5-15mg CSCI/24 hour (terminal phase)
 - Use with caution in the elderly
4. Consider low-dose diuretics if there's fluid overload

Nursing and Supportive Care (B)

1. Manage environment
 - General calm in voice & actions
 - Open windows, facilitate air movement, avoid crowding, loosen clothes

¹ Verberkt CA, JMGA, et al. Respiratory adverse effects of opioids for breathlessness: a systematic review and meta-analysis. Eur Respir J 2017;50. doi:10.1183/13993003.01153-2017

NCG Palliative Care Guidelines - Breathlessness

- Oxygen: Limit SpO₂ monitoring to intermittent use if at all - Use guided by patient comfort & not by sPO₂. A trial with a fan advisable, before starting oxygen. Disadvantages – sense of suffocation & discomfort to patient, constant source for anxiety to family, alarms
 - Ambulatory O₂ – a very selective decision considering bed-bound /limiting status vs. significant logistical disadvantages and safety (fire) risks. May be used as short bursts around exercise.
2. Communication
- Prognosis – ensure consistency within the Team
 - i. Poor prognosis - : mMRC Gr. 4-5²; FEV₁ < 30%predicted, right heart failure, poor nutrition, previous ICU/NIV/, resistant to Antibiotics, steroids trials in the past year
 - ii. Uncontrolled comorbidities, multi-system failure
 - Goals of interventions - in terms of beneficence (not effects on parameters- sPO₂) and irreversibility of vcertain contributors
 - Revisit Advance Care-Directive ACD (withhold/ withdraw mechanical ventilation). Include patient and family preferences
 - Preferred place for terminal phase, religious/ cultural needs
 - Discuss the option for of sedation for symptom relief - explain loss to communication
3. Non-invasive Ventilation (NIV - CPAP & BiPAP)
- Most benefit seen after the first hour of treatment and in hypercapnic patients
 - May be used while awaiting decisions on benefits / futility of invasive aggressive interventions
 - At the terminal phase – the use is guided entirely by patient comfort
4. Non-Pharmacological Interventions - Engage & activate care-inputs from Paramedical MDT in suitable patients
- Occupational therapy with Energy conservation techniques -activity pacing, prioritisation, aids to help functionality
 - Physiotherapy – walking aids, breath re-training, reduce work of breathing, encourage relaxation, expectorating secretions, re-conditioning,
 - Coping strategies to identify breathlessness early, improve breathing control.
 - Cognitive Behaviour therapy to identify triggers, respond mindfully, sets goals & use anxiety tool-kit
 - Pulmonary rehabilitation. Discontinue intravenous fluids.
5. Specific to terminal stages
- Accept intractability of breathlessness; Respect dignity, preferences & comfort of the patient and withhold, futile interventions that add to the distress and prolong the dying.
 - Reduce excessive secretions – Oral hygiene, position, Atropine, Glycopyrrolate;
 - Avoid Suction – restrict to oral suction

² https://www.researchgate.net/figure/The-modified-Medical-Research-Council-dyspnea-scale-MMRC-35_tbl1_23938183

NCG Palliative Care Guidelines - Breathlessness

- Institute and adjust sedation aimed at symptom relief- take informed consent from family and follow institutional guidelines (on-going analgesia with Morphine IS NOT sedation)